



Quantitative Palmar Dermatoglyphic Patterns in Cases of Idiopathic Generalized Epilepsy

Authors

Dr Kamil Khan¹, Dr Kishwor Bhandari², Dr Mohd Tabrez Alam³

Mrs. Sanju Acharya⁴, Dr Mohd Imran⁵

^{1,2,3}Assistant Professor, Dept of Anatomy, Hind Institute of Medical Sciences, Sitapur, UP, India

⁴Assistant Lecturer, Dept of Physiology, Hind Institute of Medical Sciences, Sitapur, UP, India

⁵Assistant Professor, Dept of Anatomy, JN Medical College, Aligarh Muslim University Aligarh, UP, India

Corresponding Author

Dr Kamil Khan

Assistant Professor, Department of Anatomy, Hind Institute of Medical Sciences, Sitapur, UP, India

Email: dr.kamilkhan@gmail.com, Phone: (+91) 9690125881, (+91) 9129647579

ABSTRACT

Introduction: Genetic etiology has been proposed for both idiopathic epilepsy and dermatoglyphics. Hence, the present study has been undertaken to find out the existence of any correlation between dermatoglyphics and idiopathic generalized epilepsy.

Materials and Methods: The study was conducted in the Department of Anatomy and Department of Neurology, Himalayan Institute of medical Sciences, Swami Ram Nagar, Dehradun. Total Finger Ridge Count (TFRC), Absolute Finger Ridge Count (AFRC), Angles; 'atd', 'adt' and 'dat' were analysed by utilizing finger and palmar prints.

Aim and Objectives: To analyse dermatoglyphic patterns of hands in idiopathic generalized epilepsy.

Results: TFRC and AFRC were significantly decreased when both the hands were taken together. A non-significant difference in angles ('atd', 'adt' & 'dat') was noted in both the hands.

Conclusion: Decrease in the TFRC and AFRC in persons may be a high risk of idiopathic generalized epilepsy.

Keywords: Dermatoglyphics, idiopathic generalized epilepsy, TFRC, AFRC angles ('atd', 'adt' & 'dat')

INTRODUCTION

Dr. Cummins, the father of American fingerprint analysis, coined the term "Dermatoglyphics". The term dermatoglyphics is derived from the ancient Greek words 'derma' means skin and 'glyph' meaning carving. Dermatoglyphics is the scientific study of patterns of epidermal ridges present on fingers, palms, toes and soles.¹

Although Cummins and Midlo coined the term dermatoglyphics, dermatoglyphics as a scientific discipline began with the publication of Purkinje's thesis and Galton's classic book "Fingerprints".² Dermatoglyphic patterns are constant and individualistic. Abnormalities in the epidermal ridges may result from genetic alterations occurring around the first trimester, during the

period of organogenesis and between the 13th to 60th days after fertilization.³

Epilepsy is a major health problem in developing countries including India. India has the second largest population in the world with almost more than one billion inhabitants and 70% of them living in rural communities.⁴

According to World Health Organization (WHO) and the International League Against Epilepsy (ILAE), epilepsy has been defined as a recurrent paroxysmal disorder of cerebral function characterized by sudden brief attacks of altered consciousness, motor activity, sensory phenomenon, or inappropriate behaviour caused by abnormal excessive discharge of cerebral neurons.⁵

The nervous system and the epidermal papillary ridges of the skin are developed from the ectoderm.⁶ The present study was carried out to identify an association between idiopathic generalized epilepsy and the dermatoglyphic patterns present in their hand.

MATERIALS AND METHODS

The present study was conducted in the Department of Anatomy, Himalayan Institute Of Medical Sciences, Swami Ram Nagar, Dehradun. Fifty confirmed cases of idiopathic generalized epilepsy were taken as cases who reported to the neuromedicine outpatient department (OPD). Prior approval of institutional ethical committee was obtained. A descriptive study was performed on 100 subjects. The study was performed on 50 normal individuals (26 males and 24 females) as control group and 50 patients taken as epilepsy group (32 males and 18 females). Thus 250 fingers & 50 hands of controls and cases each were analyzed in both right and left hands separately. The age groups of the subjects were between 15-50 years of either sex.

The control subjects were selected from the patients attending the neuromedicine OPD. Patients with history of epilepsy, convulsions or seizures in the past were excluded from the control group.

Selection of epilepsy group consists of patients clinically confirmed as cases of idiopathic generalized epilepsy who were attending the OPD. Patients with history of epilepsy in the past associated with a variety of cerebral or systemic disorders in the form of chronic infantile encephalopathy, severe mental deficiency, spastic or flaccid paraplegia or tetraplegia, cardiac, ocular or other diseases were excluded from epilepsy group. Epileptic patients with history of previous trauma over the head or meningeal infections were also excluded from epilepsy group.

Case recording forms were used to generate the data. A written informed consent was taken from all the subjects before taking the finger and palmar prints.

The materials used for the study were quick drying duplicating ink, rubber roller, inking slab-thick glass sheet, white paper with a glazed surface on one side of A3 size, pressure pad made up of foam, diluent and hand lens. Printing of the digits and palm were obtained using the ink method. Parameters analyzed by utilizing finger and palmar prints were Total Finger Ridge Count (TFRC), Absolute Finger Ridge Count (AFRC), Angles 'atd', 'adt' and 'dat'. The data were analyzed by using statistical software SPSS-22. The collected data were represented in the form of mean and standard deviation. The significance of differences between cases and controls was assessed by the Student's t-test at 5% level of significance.

RESULTS

Table 1: Comparison of Total Finger Ridge Count (TFRC) Between Control and Epilepsy group

Parameters		Control (n=50)	Epilepsy (n=50)	p value
Right Hand	Mean	72.38	70.54	0.19
	SD	19.58	19.06	
	SEM	2.77	2.696	
Left Hand	Mean	75.28	68.98	0.42
	SD	16.77	21.92	
	SEM	2.372	4.82	
Total	Mean	147.66	139.52	0.049
	SD	36.35	40.98	
	SEM	5.142	7.516	

Table 2: Comparison of Absolute Finger Ridge Count (AFRC) Between Control and Epilepsy Group

Parameters		Control (n=50)	Epilepsy (n=50)	p value
Right Hand	Mean	113.36	101.62	0.18
	SD	45.77	40.9	
	SEM	6.47	5.78	
Left Hand	Mean	109.36	100.06	0.31
	SD	44.69	46.04	
	SEM	6.32	6.51	
Total	Mean	222.72	201.68	0.041
	SD	90.46	86.94	
	SEM	12.79	12.29	

Table 3 a: Comparison of Angle 'atd' between Control and Epilepsy Group

Parameters		Control (n=50)	Epilepsy (n=50)	p value
Right Hand	Mean	41.6	42.04	0.14
	SD	4.5	3.79	
	SEM	0.636	0.535	
Left Hand	Mean	42.56	42.14	0.4
	SD	5.39	4.47	
	SEM	0.762	0.632	

Table 3 b: Comparison of angle ‘dat’ between Control and Epilepsy Group

Parameters		Control (n=50)	Epilepsy (n=50)	p value
Right Hand	Mean	59.38	58.82	0.4
	SD	6.44	4.72	
	SEM	0.911	0.667	
Left Hand	Mean	58.34	59.38	0.3
	SD	4.29	6.12	
	SEM	0.607	0.865	

Table 3 c: Comparison of angle ‘adt’ between Control and Epilepsy Group

Parameters		Control (n=50)	Epilepsy (n=50)	p value
Right Hand	Mean	79.1	79.66	0.6
	SD	5	4.525	
	SEM	0.848	0.64	
Left Hand	Mean	78.72	78.58	0.91
	SD	5.718	6.854	
	SEM	0.809	0.969	



Fig. 1a: Palmar Print of Left Hand of Male Control showing <adt (80), <dat (60) and <atd (40).



Fig. 1b: Palmar Print of Left hand of Male Case showing <adt (65), <dat (75) and <atd (40).

The data obtained were compiled and tabulated. Table 1 showed that there was a non significant decrease in TFRC in cases when taken for right and left hands separately (p values were 0.19 and 0.42 respectively). However, when TFRC was compared between control and epilepsy group for both hands taken together the p value was 0.049 suggesting a significant decrease in the mean value of TFRC.

Table 2 showed that there was a non significant decrease in AFRC in cases when taken for right and left hands separately (p values were 0.18 and 0.31 respectively). However, when AFRC was compared between controls and epilepsy group for both hands taken together the p value was 0.041 suggesting a significant decrease in the mean value of AFRC.

There was a non-significant increase in angle ‘atd’ and angle ‘adt’ in right hands (p >0.05) and non-significant decrease in the angles in left hands (p >0.05) of cases when compared with respective

hands of controls. However, angle 'dat' showed a non-significant decrease in right hand and a non-significant increase in the angle in left hand of epilepsy group when compared with respective hands of control group. (Fig1a & 1b).

DISCUSSION

In the present study there was a significant decrease in the mean values of TFRC (139.52 ± 40.98) and AFRC (201.68 ± 86.94) in cases when compared with both hands of controls taken together. There was a non-significant decrease in TFRC (70.54 ± 19.06) and AFRC (101.62 ± 40.9) in right hand fingers of cases when compared with their respective controls having TFRC (75.28 ± 16.77) and AFRC (113.36 ± 45.77). There was also a non significant decrease in TFRC (68.98 ± 21.92) and AFRC (100.06 ± 46.04) in left hand fingers of cases when compared with their respective controls having TFRC (72.38 ± 19.58) and AFRC (109.36 ± 44.69) (Tables 1&2).

Shawky et al., found significantly reduced TFRC in both the hands of epileptic patients in the Egyptian population.⁷ Mattos- Fiore MAB and Saldanha PH observed similar decrease in TFRC in female epileptic patients but increased values in cases of male epileptics.⁸

Contradictory findings were observed by many scientists. Mulage SK found increased TFRC and AFRC in male and female epileptics.⁹ Lal & Sureka and Filho et al., found increased incidence of TFRC and AFRC in epileptic patients.^{10,11} Rivera did not observe any difference in TFRC and AFRC of epileptic patients when compared with their controls.¹²

In the present study there was an increase in the mean value of angle 'atd' in right hand (42.04 ± 3.79)⁰ while there was a decrease in mean value of the angle in left hand (42.14 ± 4.47)⁰ of epilepsy group when compared with their respective controls i.e. 41.6 ± 4.5 ⁰ and 42.56 ± 5.39 ⁰ respectively ($p > 0.05$) (Figs. 1a & 1b). On observing the angle 'dat', a decrease in right hand (58.82 ± 4.72)⁰ and an increase in left hand

(59.38 ± 6.12)⁰ of cases were found when compared with their respective controls i.e. 59.38 ± 6.44 ⁰ and 58.34 ± 4.29 ⁰ respectively ($p > 0.05$) (Fig. 1a & 1b). Measurement in angle 'adt' revealed increased mean value of the angle in right hand (79.66 ± 4.53)⁰ but a decrease in the value in left hand (78.58 ± 6.85)⁰ of cases when compared with their respective controls i.e. 79.10 ± 5.00 ⁰ and 78.72 ± 5.72 ⁰ respectively ($p > 0.05$) (Figs. 1a & 1b).

Ranganath et al., observed an increase in the mean values of angles 'atd' and 'dat' but a decrease values of angle 'adt' in both the hands of male and female epileptic patients.⁵ Lal and Sureka noticed a decreased mean value of angles 'atd' and 'dat' but an increase in the value of angle 'adt' in right and left hands of epileptic patients.¹⁰ Goshi RC et al concluded a significant increase in 'dat' angle in the right and decrease in 'adt' angle in both hands of male epileptics.¹³ In the study conducted by Filho et al., reference has been made to the study of Rosner et al., who observed higher values of angle 'atd' in left hand of male epileptics but smaller values of the angle in both hands of female epileptics when compared with their respective controls. In the same study, Ravazi found no statistical differences in angle 'atd' in both hands and genders of epileptics when compared with their respective controls.¹¹

CONCLUSION

Decrease in the TFRC and AFRC in persons may be a high risk of idiopathic generalized epilepsy. Therefore, these parameters of significantly decreased TFRC and AFRC may be of immense help not only in identifying pre-epileptics but can be helpful in preventing serious complications of epilepsy, which is a leading problem of the world.

ACKNOWLEDGEMENTS

The authors would like to thank Dept. of Neurology, Himalayan Institute Hospital Trust, Jolly Grant, Dehradun for their guidance and support.

Funding: No funding sources

Conflict of Interest: None declared

REFERENCES

1. Dermatoglyphics. <http://en.wikipedia.org/wiki/Dermatoglyphics>.
2. Cummins H, Midlo C. Palmar and plantar epidermal ridge configurations in European Americans. *Am J Phys Anthropol*. 1926;9:471-502.
3. Kumbnani HK. Anthropologist Palmar Main Line Index - A New Methodological Approach. *Anthropologist*. 2004; 6 (2): 151-2.
4. Saha SP, Bhattacharya S, Roy BK, Basu A, Roy T, Maity B et al. A prospective incidence study of epilepsy in a rural community of West Bengal, India. *Neurology Asia*. 2008; 13:41-8.
5. Epilepsy. <http://www.who.int/mediacentre/factsheets/fs999/en/>.
6. Skin Anatomy. <http://www.enchantedlearning.com/subjects/anatomy/skin/>.
7. Shawky RM, El-Sawy M, Hashad H. Dermatoglyphic study in cases of idiopathic epilepsy. *Egyptian J Med Sci*. 1990a; 58(2):207-14.
8. Mattos-Fiore MAB, Saldanha, PH. Dermatoglyphics in juvenile epilepsy. I. Finger patterns and ridge counts. *Brazilian Journal of Genetics*. 1996; 19(1); 151-64.
9. Mulage SK. Study of palmar dermatoglyphic pattern in epileptic patients of KLES dr. Prabhakar kore hospital and medical research centre, Belgaum - a cross sectional study. 2009.
10. Lal N, Sureka RK. A Study of Dermatoglyphic Patterns in Epileptic Patients. *Journal of*
11. Filho G, Schaumann K, Kimura P, Kharitonov RA, Kozlova AI. Dermatoglyphic traits in the diagnosis and prognosis of epilepsy. *Zh Nevropatol Psikhiatr Im S S Korsakova*. 1985; 85(6):861-7.
12. Rivera C, Roman J, Arias C, Bertha A, Alvarez M. Palmer dermatoglyphics in children with febrile seizures/palmodigital dermatoglyphic phenotypes in children with febrile convulsions. *Rev Mex Pediatrics*. 1996; 63(4): 174-7.
13. Ghoshi RC, Ranganath P, Kulkarni RN, Srinivasa R. Qualitative Dermatoglyphics in Idiopathic Epilepsy. *Annals of Indian Academy of Neurology* 2004; 17: 319-2