



## Research Article

# Role of Electroconvulsive Therapy in Patients with Severe Depression

Authors

**Nikhil S Gupta<sup>1\*</sup>, Nilesh M Naphade<sup>2</sup>, Jyoti V Shetty<sup>3</sup>**

<sup>1</sup>Resident, <sup>2</sup>Professor, <sup>3</sup>Professor & Head

Department of Psychiatry, Bharati Vidyapeeth Deemed University Medical College & Research Hospital,  
Pune-411043

Corresponding Author

**Dr Nikhil Gupta**

Resident, Department of Psychiatry, Bharati Vidyapeeth Deemed University Medical College

Email: [drnikhilsgupta@yahoo.com](mailto:drnikhilsgupta@yahoo.com), Mobile: +91- 7350414881

## ABSTRACT

**Introduction:** Depression is the leading global cause of disability and approximately, 350 million people suffer from depression worldwide. Despite the availability of numerous psychopharmacological treatments, evidence indicates that only 60–70% of persons who tolerate antidepressants will respond to first-line drug therapy for major depressive disorder. Electroconvulsive therapy (ECT) is a procedure which consists of the induction of convulsive crises by means of applying an electric current through the brain for therapeutic purposes. Resistance to the antidepressant medication is the main indication for ECT. The present study was planned to evaluate changes in depressive symptoms before ECT treatment, immediately after ECT treatment and at one month after ECT treatment.

**Materials & Methods:** A total of 30 consecutive patients diagnosed with Severe Depressive Episode as per International Classification of Diseases (ICD-10) Diagnostic Criteria for Research and satisfying the eligibility criteria were taken in the study after informed consent. ECT was administered as per prevailing guidelines by means of a medical model BPE – 2000 giving bidirectional square wave pulse at a frequency of 20-90Hz. Baseline evaluation was done one day prior to ECT treatment while Post-treatment evaluation was carried out at the end of ECT treatment and at one month after ECT treatment.

**Results:** The mean age of the study subjects was 38.53 years with 66.7% females and 33.3% males. Immediately post-treatment 6 participants had HAM-D scores in the range indicating no depression (score < 8), 17 had scores in the mild to moderate depressed range (scores 8-18), 6 demonstrated severe depression (19-22), and 1 had very severe depression > 22. At 1-month follow-up, 4 patients fell in the range indicating no depressive symptoms, 19 in the mild to moderate range, 5 in the severe range, and 2 in the very severe depressed range. Mean Hamilton score was 40.03, 14.32 and 14.89, before ECT, immediately after ECT and at 1 month follow up respectively ( $p < 0.01$  and  $p = 0.19$ ).

**Conclusion:** The observation made in present study strongly suggest the benefits of ECT in reducing and providing sustained relief from the acute symptoms of depression. The biggest challenge however is to deal with stigma, prejudice and resistance to the use of such an efficient treatment.

**Keywords:** Electroconvulsive Therapy, Hamilton Rating Scale for Depression, Severe Depression.

## INTRODUCTION

Depression is the leading global cause of disability and approximately, 350 million people suffer from depression worldwide <sup>[1]</sup>. About 13% of men and 21% of women will be affected in their lifetime and, according to WHO, Major Depression will be the world's second most debilitating disease by 2020 <sup>[1]</sup>. India has about 16.6 million cases of Major Depression per year and 10.3 million cases at any time <sup>[2]</sup>.

Despite the availability of numerous psychopharmacological treatments, evidence indicates that only 60–70% of persons who tolerate antidepressants will respond to first-line drug therapy for major depressive disorder (MDD) <sup>[3]</sup>. Furthermore, at least one-third of persons with MDD who receive drug therapy will become treatment resistant <sup>[4]</sup>. Patients who do not have their symptoms fully remitted show higher recurrence risk, anxiety disorders, chronic medical illnesses, have worse psychosocial functioning, worse quality of life (QOL) and higher number of symptoms during follow-up <sup>[5-7]</sup>.

Electroconvulsive therapy (ECT) is a procedure which consists of the induction of convulsive crises by means of applying an electric current through the brain for therapeutic purposes. Resistance to the antidepressant medication is the main indication for ECT <sup>[8-11]</sup>. Besides, ECT provides a faster response as compared to medication, which is necessary in severe situations that need emergency intervention, such as catatonia and suicide risk <sup>[12]</sup>. The American Psychiatric Association has already included these situations as indication of ECT as the first choice <sup>[13]</sup>. The efficacy of ECT to treat depressive symptoms has been established by means of innumerable studies developed during the last decades <sup>[14]</sup>. According to Prudic, ECT is the most effective biological treatment currently available for depression, and no other treatment up to now has shown to be superior to ECT in the treatment of major depression in controlled studies <sup>[15]</sup>.

Not much research has been done in this area in our country. The proposed study was thus planned

to evaluate changes in Depressive symptoms before ECT treatment, immediately after ECT treatment and at one month after ECT treatment.

## MATERIALS AND METHODS

A Prospective Interventional study was conducted in patients with Severe Depression receiving Electroconvulsive Therapy during Oct' 2015 to Sept' 16. A total of 30 consecutive patients diagnosed with Severe Depressive Episode as per International Classification of Diseases (ICD-10) Diagnostic Criteria for Research and satisfying the eligibility criteria were taken in the study after informed consent <sup>[16]</sup>.

### Inclusion Criteria

- All patients above 18 years of age.
- All OPD and IPD patients diagnosed with Severe Depressive Episode (as per ICD-10) receiving Electroconvulsive Therapy.
- All patients willing to participate in the study by means of Informed consent.
- All patients giving Informed consent to undergo Electroconvulsive Therapy prior to every treatment.

### Exclusion Criteria

- All patients who had received Electroconvulsive Therapy within past one year.
- All individuals with Neurological deficits, Serious medical illnesses and Mental Retardation.
- All individuals with other Psychiatric and Neurological disorders like Dementia, Mania, Psychosis, and Substance use disorders.

### Methodology

A written Informed Consent after explaining the procedure in detail was taken before each ECT treatment. Baseline evaluation was done one day prior to ECT treatment while Post-treatment evaluation was carried out at the end of ECT treatment and at one month after ECT treatment. All the observations were recorded on a specially designed proforma and validated by two experienced psychiatrists independently.

### Depression assessment

The Hamilton Rating Scale for Depression (HAM-D) <sup>[17]</sup> was used to assess depressive symptomatology in all participants at pre-treatment, post-treatment, and follow-up testing sessions. This study utilized the 21-item interview which rates the severity of symptoms observed in depression such as low mood, insomnia, agitation, anxiety and weight loss. The questionnaire is one of the most commonly used scales for rating depression in medical research. The investigator rates the patient by interviewing them and observing the patient's symptoms. Each question has from three to five possible responses that increase in severity. The total score represents a sum of the values for the questions. Scores ranging 0-7 were normal, from 8 to 13 indicate mild depression symptomatology, 14 to 18 indicate moderate depression, 19 to 22 indicate severe depression and scores above 22 are interpreted as indicating very severe depression. The maximum score one can receive on this measure is 66.

### Electroconvulsive Therapy

ECT was administered as per prevailing guidelines by means of a medical model BPE – 2000 giving bidirectional square wave pulse at a frequency of 20-90Hz <sup>[18]</sup>. Seizure threshold was determined by the half age method of Petride and Fink <sup>[19]</sup>.

### Statistical Analysis

All the collected data was entered in Microsoft Excel Sheet 2007. The data was then transferred and analyzed using SPSS ver. 17. Qualitative data was represented in the form of frequency and percentage and analyzed by chi-square test while quantitative data was represented using mean +/- S.D and analyzed using Wilcoxon Sign Rank Test. A p-value of <0.05 was taken as level of significance.

### RESULTS

The mean age of the study subjects was 38.53 years with 66.7% females and 33.3% males.

Family History of Psychiatric Illness was seen in 76.7% subjects. Most of the subjects had between 2-4 episodes during their course of illness. Other psychiatric morbidities were seen in 40% of the cases (Table 1). Immediately post-treatment 6 participants had HAM-D scores in the range indicating no depression (score < 8), 17 had scores in the mild to moderate depressed range (scores 8-18), 7 demonstrated severe/ very severe depression ( $\geq 19$ ). These data indicate that the majority of the participants responded to ECT treatment. At 1-month follow-up, 4 patients fell in the range indicating no depressive symptoms, 24 in the mild to moderate range and 2 in the severe/ very severe range. Mean Hamilton score was 40.03, 14.32 and 14.89, before ECT, immediately after ECT and at 1 month follow up respectively ( $p < 0.01$  and  $p = 0.19$ ) (Table 2 and 3).

**Table 1.** Distribution of patients according Baseline Characteristics

Baseline Characteristics (n=30)		
Variables	N	%
Age group (yrs)		
$\leq 30$	2	6.7%
31-40	16	53.3%
41-50	11	36.7%
$> 50$	1	3.3%
Sex		
Female	20	66.7%
Male	10	33.3%
Family History of Psychiatric Illness		
No	7	23.3%
Yes	23	76.7%
Course of Present Illness		
Gradual	22	73.3%
Sudden	8	26.7%
No. of Episodes		
1	4	13.3%
2-4	25	83.3%
$> 4$	1	3.3%
Psychiatric Co-morbidity		
No	18	60.0%
Yes	12	40.0%

**Table 2.** Distribution of subjects as per severity of depression

Hamilton Score	Baseline		Immediate After ECT		1 month F/u	
	N	%	N	%	N	%
No Depression	0	0.0%	6	20.0%	4	13.3%
Mild	0	0.0%	17	56.7%	19	63.3%
Moderate	0	0.0%	6	20.0%	5	16.7%
Severe	30	100.0%	1	3.3%	2	6.7%
Total	30	100.0%	30	100.0%	30	100.0%
p- value	< 0.01				0.71	

**Table 3.** Distribution of subjects based on mean change in Hamilton score

Hamilton Scale	Mean	SD	p- value
Baseline	40.03	8.76	<0.01
Immediately after ECT	14.32	9.62	
1 month After ECT	14.89	9.54	0.19

## DISCUSSION

Among the most vexing predicaments for many clinicians is providing dispassionate guidance regarding electroconvulsive therapy (ECT) to a patient and family members of a severely impaired, suicidal, medication-resistant patient, who has benefited marginally from psychotherapy. The American Psychiatric Association (2001) regards it as a very effective option when other treatments fail <sup>[1]</sup>. Present hospital based prospective observational study was conducted to evaluate changes in depressive symptoms and their sustenance in patients with severe depression.

In present study, all the 30 subjects had severe/very severe depression before ECT therapy. Immediately post-treatment 6 participants had HAM-D scores in the range indicating no depression, 17 had scores in the mild to moderate depressed range, 7 demonstrated severe/ very severe depression. Our findings indicate that the majority of the participants responded to ECT treatment. At 1-month follow-up, 4 patients fell in the range indicating no depressive symptoms, 24 in the mild to moderate range and 2 in the severe/ very severe range. Mean Hamilton score was 40.03, 14.32 and 14.89, before ECT, immediately

after ECT and at 1 month follow up respectively ( $p < 0.01$  and  $p = 0.19$ ). These findings shows that ECT provided sustained relief from symptoms of depression.

In results of the study by Saccoman et al. <sup>[20]</sup> indicate that ECT is an effective treatment of severe depression. Their study showed that 88% of the participants responded immediately after completion of a standard series of bifrontal ECT. Our success rate is also comparable to the 70 to 90% response rate for all types of ECT reported by the American Psychiatric Association's (2010) practice guidelines regarding treatment for depression <sup>[21]</sup>, and higher than remission rates of 55% and 64% found in multisite research consortium projects <sup>[22,23]</sup>.

Like other investigations, the current study also found that few participants experienced a return of depressive symptoms 1 month after treatment cessation and that on average participants demonstrated an increasing trend in depression symptomatology from post-treatment to follow-up. Specifically, 6.5% of the participants who were classified as 'no depression' as per Hamilton scale were later classified as 'mild depression' after the cessation of treatment. Saccoman et al. <sup>[20]</sup> found the relapse rate of 13.5% in their study.

Relapse rate in our study is very low as compared to some other studies of ECT (such as Prudic et al. [15], who reported a 64% 24-week relapse rate in their 2004 publication and Grunhaus et al. [24], who reported a 28.5% 3-month relapse rate in their 2001 study).

Very low relapse rates at follow-up in our study may be due to the lack of a longer follow-up period. This is considered a limitation of this study. The present data are restricted to only 1 month follow-up which is much shorter than the follow-up periods of 24 weeks from Prudic et al. [15] and 3 months in Grunhaus et al. [24] This provides reason to be cautious in making strong conclusions about ECT relapse rates. It is plausible that given the trend in the current sample of rising symptomatology longer follow-up periods may have resulted in a higher relapse rate. But still, ECT can be considered an effective course of action for severe, medication-resistant, depressive conditions.

## CONCLUSION

The observation made in present study strongly suggest the benefits of ECT in reducing and providing sustained relief from the acute symptoms of depression. The biggest challenge however is to deal with stigma, prejudice and resistance to the use of such an efficient treatment.

## REFERENCES

1. The World Health Organization. The global burden of disease: 2004 update, table A2: burden of disease in DALYs by cause, sex and income group in WHO regions, estimates for 2004. Geneva, Switzerland: WHO, 2008. [http://www.who.int/healthinfo/global\\_burden\\_disease/GBD\\_report\\_2004update\\_AnnexA.pdf](http://www.who.int/healthinfo/global_burden_disease/GBD_report_2004update_AnnexA.pdf) (accessed 6 Mar 2014)
2. S Saxena, ADhawan. Disability and Burden of depressive disorders. The National Medical Journal of India; vol.12, No.2, 1999
3. Souery D, Papakostas GI, Trivedi MH. Treatment resistant depression. J Clin Psychiatry 2006;67(Suppl 6):16–22.
4. Fava M, Davidson KG. Definition and epidemiology of treatment resistant depression. Psychiatr Clin North Am 1996;19:179–200.
5. Kennedy N, Foy K. The impact of residual symptoms on outcome of major depression. Curr Psychiatry Rep. 2005;7(6):441-6.
6. Rush AJ, Kraemer HC, Sackeim HA, Fava M, Trivedi MH, Frank E, Ninan PT, Thase ME, Gelenberg AJ, Kupfer DJ, Regier DA, Rosenbaum JF, Ray O, Schatzberg AF, ACNP Task Force. Report by the ACNP task force on response and remission in major depressive disorder. Neuropsychopharmacology. 2006;31(9):1841-53.
7. Viinamaki H, Hintikka J, Tolmunen T, Honkalampi K, Haatainen K, Koivumaa-Honkanen H. Partial remission indicates poor functioning and a high level of psychiatric symptoms: a 3-phase 6-year follow-up study on major depression. Nord J Psychiatry. 2008;62(6):437-43.
8. Husain SS, Kevan IM, Linnell R, Scott AI. Electroconvulsive therapy in depressive illness that has not responded to drug treatment. J Affect Disord. 2004;83(2-3):121-6.
9. Freeman C, Hendry J, Fergusson G. National audit of electroconvulsive therapy in Scotland; 2000. Available from: <http://www.sean.org.uk>.
10. Braga RJ, Petrides G. Somatic therapies for treatment-resistant psychiatric disorders. Rev Bras Psiquiatr. 2007;29(Suppl 2):S77-84.
11. Husain SS, Kevan IM, Linnell R, Scott AI. What do psychiatrists mean by medication resistance as an indication for electroconvulsive therapy? J ECT. 2005;21(4):211-3.

12. McCall WV. What Does Star\*D Tell Us About ECT? J ECT. 2007;23(1):1-2.
13. American Psychiatric Association Task Force on Electroconvulsive Therapy. The practice of electroconvulsive therapy: recommendations for treatment, training and privileging. Washington, DC: American Psychiatric Association; 2001.
14. Abrams R. Electroconvulsive therapy. 4th edn. Oxford, UK: Oxford University Press, 2002.
15. Prudic J. Electroconvulsive therapy. In: Sadock BJ, Sadock VA, editors. Kaplan & Sadock's comprehensive textbook of psychiatry. 8th ed. Philadelphia: Lippincott Williams & Wilkins; 2005. p.2968-83.
16. World Health Organization. The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines. Geneva: World Health Organization; 1992.
17. Hamilton M. A rating scale for depression. Journal of Neurology, Neurosurgery & Psychiatry. 1960 Feb 1;23(1):56-62.
18. Davidson JR. Major depressive disorder treatment guidelines in America and Europe. The Journal of clinical psychiatry. 2009 Dec;71:04-17.
19. Petrides G, Fink M. The " half-age" stimulation strategy for ECT dosing. The Journal of ECT. 1996 Sep 1;12(3):138-46.
20. Saccoman ML. An investigation of the cognitive effects of electroconvulsive therapy in the treatment of severe depression (Doctoral dissertation, The University of Utah).
21. American Psychiatric Association. (2010). Practice guideline for the treatment of patients with major depressive disorder (3rd Ed.) American Psychiatric Association: Arlington, VA.
22. Kellner CH, Knapp RG, Petrides G, et al. Continuation electroconvulsive therapy vs pharmacotherapy for relapse prevention in major depression: a multisite study from the Consortium for Research in Electroconvulsive Therapy (CORE). Arch Gen Psychiatry 2006; 63:1337.
23. Sackeim HA, Dillingham EM, Prudic J, et al. Effect of concomitant pharmacotherapy on electroconvulsive therapy outcomes: short-term efficacy and adverse effects. Arch Gen Psychiatry 2009; 66:729.
24. Grunhaus L, Shipley JE, Eiser A, Pande AC, Tandon R, Remen A, Greden JF. Shortened REM latency PostECT is associated with rapid recurrence of depressive symptomatology. Biological psychiatry. 1994 Aug 15;36(4):214-22.