



## Methicillin-Resistant *Staphylococcus aureus* Screening in Elective Surgery Patients in CCMMC Hospital Durg

Authors

**Dr Subhasis Mandal<sup>1</sup>, Dr Rolly Kaushik<sup>2</sup>**

<sup>1</sup>Assistant Professor Department of Surgery CCM Medical College, Kachandur, Durg (CG)

<sup>2</sup>Assistant Professor Department of Anatomy CCM Medical College, Kachandur, Durg (CG)

Corresponding Author

**Dr Rolly Kaushik**

Assistant Professor Dept. of Anatomy, CCM Medical College Kachandur, Durg (CG), India

### Abstract

**Background:** *Surgical site infection (SSI) is a major contributor of healthcare associated infection. Staphylococcus aureus is a pathogen responsible for both community-acquired as well as hospital-associated infections. SSI is associated with substantial increased of postoperative hospital stay, rates of hospital readmission, hospital cost, functional disability, and mortality rate. Patients in hospitals which are infected and colonized mediate the dissemination of (Methicillin Resistant Staphylococcus aureus) MRSA strains. Therefore it is necessary to know the prevalence of MRSA and its antibiotic sensitivity pattern for proper treatment of the patients and thereby reducing the morbidity and mortality of the patients.*

**Methods:** *The cross sectional study was conducted in CCMMC Durg. All adult patients admitted to the hospital and assigned for an elective surgery were chosen to collect data on MRSA. Swabs were taken from the anterior nares and axilla. Antibiotic susceptibility testing was performed by the disk diffusion method using guidelines established by the Clinical and Laboratory Standards Institute (CLSI) and MRSA was detected by testing with Cefoxitin disc*

**Results:** *A total of 121 patients (42.6%) were colonized with Staphylococcus epidermidis and 59 patients (20.77%) had positive nasal swabs for Methicillin-Sensitive Staphylococcus aureus (MSSA) and 24 (8.45%) patients were Methicillin-Resistant Staphylococcus aureus (MSSA).*

**Conclusion:** *MRSA strains are more prevalent in tertiary care hospital and multidrug resistant MRSA prevalence is higher in surgical units. Selective swabbing and decolonization for high risk preoperative patients may be more appropriate. The most effective way to prevent MRSA is surveillance of antibiotic resistance profiles of Staphylococcus aureus and there should be formulation of antibiotic policy in the hospital. Thus, control of MRSA is essential to curtail the introduction and spread of infection in the hospitals. This can be achieved by using universal precautions and conducting regular epidemiological studies.*

### Introduction

Surgical site infection (SSI) remains a major contributor of healthcare associated infection

(HAI). Based on Centers for Disease Control and Prevention (CDC) prevalence survey in 2011.<sup>1</sup> *Staphylococcus aureus* is a pathogen responsible

for both community-acquired as well as hospital-associated infections.

*Staphylococcus aureus* remains the leading cause of SSI, with half of the *S. Aureus* were found resistant to methicillin<sup>2</sup>. *Staphylococcus aureus* is responsible for causing a variety of human infections, which range from minor skin diseases to life-threatening infections.<sup>3</sup> Methicillin resistant *Staphylococcus aureus* (MRSA) were reported in October 1960.<sup>4</sup> SSI was associated with substantial increased of postoperative hospital stay, rates of hospital readmission, functional disability, hospital cost, and mortality rate<sup>5</sup>

Routine decolonization of MRSA with chlorhexidine bathing and mupirocin nares application before surgery is becoming an interesting strategy option to reduce number of SSI.<sup>6,7</sup>. The emergence of chlorhexidine-resistant bacteria and mupirocin resistance are two concerns raised with the wide spread application of this strategy. Active screening followed by selective decolonization is another strategy but with associated with relatively higher cost due to additional diagnostic expense.<sup>8</sup>

Over the last two or three decades, methicillin resistant *S. aureus* (MRSA) strains have become endemic in hospitals worldwide and it is now incipient community pathogen in many geographical regions. Due to increased incidence of MRSA vancomycin has been used as a empirical treatment for *Staphylococcus aureus* infections which in turn results in the emergence of VRSA (vancomycin resistant *Staphylococcus aureus*).<sup>9</sup>

Patients in hospitals those who are infected and colonized mediate the dissemination of MRSA strains<sup>10</sup>, these factors increases the chance of emergence and spread of MRSA which lead to repeated hospitalization, indiscriminate use of antibiotics, intravenous drug abuse and indwelling medical devices.<sup>11</sup> Thus, control of MRSA is essential to curtail the introduction and spread of infection in the hospitals. This can be achieved by using universal precautions and conducting regular epidemiological studies.<sup>12</sup>

Therefore it is necessary to know the prevalence of MRSA and its antibiotic sensitivity pattern for proper treatment of the patients and thereby reducing the morbidity and mortality of the patients.

### Material and Methods

**Study design:** The cross sectional study was conducted in CCMMC Durg, a referral hospital with Total capacity for inpatient is 750 beds. All adult patients admitted to the hospital and assigned for an elective surgery were chosen to collect data on MRSA

**Inclusion criteria:** All patients aged  $\geq 18$  years old, assigned for an elective surgery in the next 24 h, and willing to participate in the study were included in the study. Swabs were taken from the anterior nares and axilla, which are the common colonization sites for MRSA. Written informed consent was asked before obtaining the nasal swab.

Nasal swabs were collected using a sterile dry cotton swab from all eligible patients approximately 10-12 hours before the surgery. Swabs were immediately transported to the Microbiology department for further processing

**Risk factors for MRSA:** Risk factors such as history of antibiotics usage in the past 3 months, history of hospitalization, history of catheterization, history of invasive procedures, history of referral from other hospital was obtained from the patients selected for elective procedure.

**Antimicrobial susceptibility testing:** Antibiotic susceptibility testing was performed by the disk diffusion method using guidelines established by the Clinical and Laboratory Standards Institute (CLSI) and MRSA was detected by testing with Cefoxitin disc<sup>13</sup>.

### Results

Total 284 Swabs were taken from the patients with elective surgery from Jun2015 to Dec 2017 admitted in CCMMC Durg. A total of 121 patients (42.6%) were colonized with *Staphylococcus*

epidermidis and 59 patients (20.77%) had positive nasal swabs for Methicillin-Sensitive *Staphylococcus aureus* (MSSA) and 24 (8.45%) patients were Methicillin-Resistant *Staphylococcus aureus* (MSSA).

### Discussion

*Staphylococcus aureus* is one of the major pathogen causing skin and soft tissue infections in the community as well as invasive infections in patients. Proportion of MRSA varies from country to country, in Sweden it is 0.4% while in Belgium it is as high as 48.4%.<sup>14</sup>

Prevalence of MRSA in our study period was found to be 8.45%, in nasal and skin carriage of preoperative patients. This prevalence was similar to the study conducted by Oh ML et al in Singapore<sup>15</sup>. However some reported prevalence of MRSA from 23 to 31 %<sup>16,17</sup>.

While Hadley, et al. reported prevalence of MRSA colonization anterior nasal was 3.5% among patients underwent total joint replacement in hospital in United States.<sup>18</sup> Prevalence of 4.25% was reported in retrospective study among cardiothoracic and neurological surgical patients in United States<sup>19</sup>. Difference in prevalence in hospitals shows the need of routine or selective decolonization protocol in preoperative patients.

More studies are required to adopt the policy of routine decolonization preoperatively as Routine screening to identify MRSA colonization by culture are not only cost burdening for developing countries, but also time consuming. PCR can give rapid result for screening but it is not readily available in all hospitals. Routine decolonization with mupirocin may be more cost effective however it may raise the possibility of resistance and lead to treatment failure.<sup>20</sup> Hence Selective swabbing and decolonization for high risk preoperative patients may be more appropriate for limited resources countries.

Association of multidrug resistant MRSA adds to the problem and hospital dust is more dangerous than other or roadside dust as former may contain MRSA which may multidrug resistant and may

create problem in the hospital specially in surgical units.

Researchers from other countries also observed that these MRSA isolates are becoming multidrug resistant and were susceptible only to glycopeptides antibiotics such as vancomycin. Low level resistance even to vancomycin is also emerging.<sup>21</sup>

### Conclusion

To conclude MRSA strains are more prevalent in tertiary care hospital and multidrug resistant MRSA prevalence is higher in surgical units of our hospital. Selective swabbing and decolonization for high risk preoperative patients may be more appropriate but more studies are required. The most effective way to prevent MRSA is surveillance of antibiotic resistance profiles of *Staphylococcus aureus* and there should be formulation of antibiotic policy in the hospital.

### References

1. Centers for Disease Control and Prevention. HAI data and statistics. In: Healthcare-associated infections. CDC. 20 Apr 2016.
2. Cavalcanti SM, França ER, Cabral C, Vilela MA, Montenegro F, Menezes D, Medeiros AC Braz J. Prevalence of *Staphylococcus aureus* introduced into intensive care units of a University Hospital. *Infect Dis*. 2005 Feb; 9(1):56-63.
3. Tiwari HK, Das AK, Sapkota D, Sivrajan K, Pahwa VK. Methicillin resistant *Staphylococcus aureus*: prevalence and antibiogram in a tertiary care hospital in western Nepal. *J Infect Dev Ctries*. 2009 Oct 22; 3(9):681-4.
4. Stefani S, Chung DR, Lindsay JA, Friedrich AW, Kearns AM, Westh H, Mackenzie FM. Methicillin-resistant *Staphylococcus aureus* (MRSA): global epidemiology and harmonisation of typing

- methods. *Int J Antimicrob Agents*. 2012;39(4):273–282.
5. Reichman DE, Greenberg JA. Reducing surgical site infections: a review. *Obstet Gynecol*. 2009 Fall; 2(4):212-21.
  6. Bebko SP, Green DM, Awad SS. Effect of a preoperative decontamination protocol on surgical site infections in patients undergoing elective orthopedic surgery with hardware implantation. *JAMA Surg*. 2015 May; 150(5):390-5.
  7. Kim DH, Spencer M, Davidson SM, Li L, Shaw JD, Gulczynski D, Hunter DJ, Martha JF, Miley GB, Parazin SJ, Dejoie P, Richmond JC. Institutional prescreening for detection and eradication of methicillin-resistant *Staphylococcus aureus* in patients undergoing elective orthopaedic surgery. *J Bone Joint Surg Am*. 2010 Aug 4; 92(9):1820-6.
  8. Roth VR, Longpre T, Coyle D, Suh KN, Taljaard M, Muldoon KA, Ramotar K, Forster A. Cost Analysis of Universal Screening vs. Risk Factor-Based Screening for Methicillin-Resistant *Staphylococcus aureus* (MRSA). *PLoS One*. 2016; 11(7):e0159667.
  9. Lowy FD. *Staphylococcus aureus* infections. *N Engl J Med*. 1998 Aug 20; 339(8):520-32.
  10. Rajadurai pandi K, Mani KR, Panneerselvam K, Mani M, Bhaskar M, Manikandan P. Prevalence and antimicrobial susceptibility pattern of methicillin resistant *Staphylococcus aureus*: a multicentre study. *Indian J Med Microbiol*. 2006 Jan; 24(1):34-8.
  11. Anupurba S, Sen MR, Nath G, Sharma BM, Gulati AK, Mohapatra TM. Prevalence of methicillin resistant *Staphylococcus aureus* in a tertiary referral hospital in eastern Uttar Pradesh. *Indian J Med Microbiol*. 2003 Jan-Mar; 21(1):49-51.
  12. Vidhani S, Mehndiratta PL, Mathur MD. Study of methicillin resistant *S. aureus* (MRSA) isolates from high risk patients. *Indian J Med Microbiol*. 2001 Apr-Jun; 19(2):13-6.
  13. Clinical and Laboratory Standards Institute (CLSI). M100-S25 Performance standards for antimicrobial susceptibility testing; Twenty-fifth informational supplement. Wayne: Clinical and Laboratory Standards Institute; 2015.
  14. Sader HS, Farrell DJ, Jones RN. Antimicrobial susceptibility of Gram-positive cocci isolated from skin and skin-structure infections in European medical centres. *Int J Antimicrob Agents*. 2010 Jul; 36(1):28-32.
  15. Oh ML, Tan SY. P054: Prevalence and risk factor analysis for methicillin-resistant *Staphylococcus aureus* colonization in an acute care hospital. *Antimicrob Resist Infect Control*. 2013;2:P54. doi: 10.1186/2047-2994-2-S1-P54
  16. Rajadurai pandi K, Mani KR, Panneerselvam K, Mani M, Bhaskar M, Manikandan P. Prevalence and antimicrobial susceptibility pattern of methicillin resistant *Staphylococcus aureus*: a multicentre study. *Indian J Med Microbiol*. 2006 Jan; 24(1):34-8.
  17. Majumder D, Bordoloi JS, Phukan AC, Mahanta. Antimicrobial susceptibility pattern among methicillin resistant *Staphylococcus* isolates in Assam. Majumder D, Bordoloi JS, Phukan AC, Mahanta. *Indian J Med Microbiol*. 2001 Jul-Sep; 19(3):138-40.
  18. adley S, Immerman I, Hutzler L, Slover J, Bosco J. *Staphylococcus aureus* decolonization protocol decreases surgical site infections for total joint replacement. *Arthritis*. 2010.
  19. Apoor R, Barnett CJ, Gutmann RM, Yildiz VO, Joseph NC, Stoicea N, et al. Preoperative prevalence of *Staphylococcus aureus* in cardiothoracic and neurological

surgical patients. *Front Public Health*. 2014;2:204.

20. Anderson MJ, David ML, Scholz M, Bull SJ, Morse D, Hulse-Stevens M, et al. Efficacy of skin and nasal povidone-iodine preparation against mupirocin-resistant methicillin-resistant *Staphylococcus aureus* and *S. aureus* within the anterior nares. *Antimicrob Agents Chemother*. 2015.
  21. Assadullah S, Kakru DK, Thoker MA, Bhat FA, Hussain N, Shah A. Emergence of low level vancomycin resistance in MRSA. *Indian J Med Microbiol*. 2003 Jul-Sep; 21(3):196-8.
-