



## Original Article

# Clinical Spectrum of Fever Cases in Correlation with Hemato-Biochemical Parameters- An Experience from a North Indian Hospital

Authors

**Dr Ankur Malhotra<sup>1</sup>, Dr Viney Sambyal<sup>2\*</sup>, Dr Sanjay Sharma<sup>3</sup>**

<sup>1</sup>Senior Resident Medicine, Post-Graduate Department of Medicine, GMC Jammu

<sup>2</sup>Lecturer Medicine, Post-Graduate Department of Medicine, GMC Jammu

<sup>3</sup>Lecturer Medicine, Post-Graduate Department of Medicine, GMC Jammu

\*Corresponding Author

**Dr Viney Sambyal**

Email: [vineysambyal@gmail.com](mailto:vineysambyal@gmail.com), Ph- 9419152854

## Abstract

**Introduction:** *The purpose of this study is to study the clinical spectrum of fever cases, outline the hemato-biochemical and serological parameters and delineate a syndromic approach to tropical infections in a Tertiary hospital of North India.*

**Methods:** *The study included 926 cases of fever admitted to our hospital, GMC Jammu, in medical wards over a period of six months and correlating their clinical manifestations with the lab parameters (hematological, biochemical, serology) and imaging modalities.*

**Results:** *Out of 926 cases admitted with fever, 78 (8.4 %) cases were undifferentiated viral fever, 140 (15.2%) were enteric fever, 85(9.2%) were dengue, 126(13.6%) were malaria, 28(3%) of hepatitis A and 26(2.8%) of hepatitis E, 14(1.5%) of varicella, 86(9.4%) of pneumonia and 81(8.7%) of UTI. 45(4.9%) liver abscess were reported along with 186(20.1%) cases of Rickettsial Fever, 0.2% of leptospirosis and 0.6% of Tuberculosis .*

**Conclusion:** *In our study, predominantly younger population (<55 years) was most commonly affected. The most common etiology was found to be Rickettsial Fever followed by Enteric Fever and Malaria. This upholds the necessity for better health awareness, early transport and presentation along with better diagnostic tools at primary health care facilities.*

**Keywords:** *Hepatitis, dengue fever, Enteric fever, Rickettsial Fever, Malaria.*

## Introduction

Fever without diagnosis is an unpleasant situation which neither the patient nor the treating physician would like to be in. However this is the single most baffling problem faced by every physician in their daily practice. Fever is a common reason for seeking healthcare in low-and middle-income countries<sup>[1]</sup>. Among patients with

febrile illness requiring admission case fatality ratios are high, sometimes exceeding 20%<sup>[2-6]</sup>. To describe epidemiologic patterns and to identify data gaps in our understanding of severe febrile illness in low resource areas, we sought to systematically review prospective hospital based study of the etiology of febrile illness. Every year different parts of India are hit by seasonal fevers

especially during the monsoon and post monsoon period between June to September as stated other studies<sup>[7,8]</sup>. Acute undifferentiated febrile illness can be differentiated from Fever of unknown origin by fever duration, and progression as described by Phuong et al.<sup>[9]</sup>.

### Methodology

A cross-sectional study was conducted over a period of 6 months from 1 June 2018 to 1 Dec 2018. All 15 to 85 years aged admitted cases with history of fever over 38.3 degrees C for < 3 weeks of duration without any specific localizing signs or symptoms were enrolled for the study. All critically ill patients requiring direct admission to ICU, cases of localized infection like abdominal infections, meningitis etc., patients of severe sepsis with septic shock, haematological malignancies, autoimmune disorders, those on immunosuppressant, fever > 3 weeks' time (PUO) were excluded from the study. Following information was retrieved: demographic data including age, sex and clinical hospitalization data including rash, vomiting, headache, joint pain, diarrhoea. Relevant biochemical, haematological and serological data were also retrieved for eg. Dengue NS1 Ag/IgM antibody, malarial parasite slide/card, Rickettsial serology, typhidot assay. Laboratory results were interpreted as per standard diagnostic criteria's for different etiologies.

Data thus collected were incorporated in to Microsoft excel sheet and analysed using IBM SPSS version 20 software (trial version).

### Results

A total of 1560 patients were admitted during study period of 6 months. Out of these, 926 were enrolled in study and 634 were excluded as per exclusion criteria. Of total 926 patients, males (58.3%) were more commonly affected in comparison to females (41.7%). (Table 1) Majority of the patients were young, in their second, third and fourth decades of life with 57.9% of patients between 15 to 45 years of age.

2.8% of patients were above age of 86 years. (Table 2)

Symptoms associated with fever were vomiting (44.3%), malaise (62.6%), joint pain (30.7%), cough (60 %), rashes (41.5%) and diarrhoea (15.8%). (Table 3)

On general examination 384(41.5%) patients had icterus while pallor was present in 206(22.2%) patients. 504(54.4%) cases were found to have hepatosplenomegaly. Although 410(44.3%) cases had added breath sounds.(Table 4)

On further hematological and biochemical workup, 626 (67.6%) cases had leukocytosis while 184 (19.9%) cases had leucopenia. 402 (43.4%) patients have thrombocytopenia. (Table 5)

516(55.7%) cases were found to have elevated bilirubin while deranged SGOT, SGPT and RFT were found in 480(51.8%), 386(41.7%) and 204(22%) cases respectively. (Table 6)

Among these 926 study participants, etiologically 13.6% were diagnosed as Malaria, 15.2 % as typhoid, 9.2 % as dengue, 20.1% as Rickettsial Fever, 8.7% as Urinary tract infections and 8.4% remains undiagnosed. (Table 7).

History of hypertension, diabetes, cardiovascular diseases and COPDs was asked. These all comorbidities, combined together were present in 9.1 % of patients.

Average length of stay for majority was between 3 to 5 days with mean length of stay was 4.36 days.

**Table 1** Sex Distribution

Sex	Number Of Patients (N)	Percentage (%)
Male	540	58.3
Female	386	41.7
Total	926	100

**Table 2-** Age Distribution

Age( In Yrs)	Number Of Patients (n)	Percentage (%)
15-25	164	17.7
26-35	174	18.8
36-45	198	21.4
46-55	126	13.6
56-65	78	8.4
66-75	74	8
76-85	86	9.3
>86	26	2.8
Total	926	100

**Table 3-Associated Symptoms with Fever**

Symptoms	Number Of Patients (n)	Percentage (%)
Cough	246	60
Rash	384	41.5
Burning Micturation	216	23.3
Diarrhoea	146	15.8
Rigor And Chills	410	44.3
Malaise	580	62.6
Joint Pains	284	30.7
Chest Pain	136	14.7
Breathlessness	190	20.5
Pain Abdomen	289	31.2
Vomiting	410	44.3

**Table 4 Clinical Signs in Patients**

Clinical Signs	Number of Patients (n)	Percentage (%)
Pallor	206	22.2
Icterus	384	41.5
Hepato-Splenomegaly	504	54.4
Added Breath Sounds	410	44.3
Pedal Edema	126	13.6
Neck Stiffness		
Renal Tenderness	56	6.04

**Table 5 Hematological Finding**

Hematological Finding	Number of Patients (n)	Percentage (%)
Leucocytosis (>11000/cumm)	626	67.6
Leucopenia (<4000/cumm)	184	19.9
Thrombocytopenia (<1.5LAC/cumm)	402	43.4
Abnormal PTI	101	10.9

**Table 6 Patients with Abnormal LFT and KFT**

Parameter	Number of Patients (n)	Percentage (%)
Raised Sgot	480	51.8
Raised Sgpt	386	41.7
Raised Bilirubin	516	55.7
Raised S.Urea Or Creatinine	204	22

**Table 7 Different Tropical Infections/Etiology of Fever Detected in the Study**

Etiology	Number Of Patients (N)	Percentage (%)
Enteric Fever	140	15.2
Dengue	85	9.2
Malaria	126	13.6
Pneumonia	86	9.4
UTI	81	8.7
Meningo-Encephalitis		
Rickettsial Fever	186	20.1

Hepatitis A	28	3
Hepatitis E	26	2.8
Varicella	14	1.5
Drug Fever	18	1.9
Undifferentiated Viral Fever	78	8.4
Liver Abscess	45	4.9
HIV	5	0.5
Leptospirosis	2	0.2

**Discussion**

Our study showed high incidence of tropical infections such as Clinical Enteric fever, Dengue, Malaria, Varicella, UTI and hepatitis A and E with most common infection in our area as Rickettsial Fever. In our study 85(9.2%) cases were diagnosed as dengue fever.

It is useful to remember that geriatric age group with UTI or pneumonia often present with febrile encephalopathy with few other clinical signs. In our study 86(9.4%) cases had pneumonia (clinic-radiologically proven). 186 patients presented with rickettsial disease .The disease occurs 7-10 days after the bite of larval from of trombiculid mite and present with fever, relative bradycardia, severe myalgias and a non-pruritic macular-papular rash . Splenomegaly was detected in 44 to 52% of the cases described in the recent outbreaks in the Shimla hills while the characteristic eschar was seen only in a minority of cases (9.5%)<sup>[10,11]</sup>. Both of our cases had characteristic eschar formation with splenomegaly and serositis. Clinical jaundice and deranged LFTs have been described in outbreaks of scrub typhus seen in Shimla hills with a reported incidence of jaundice in 10-52% and elevated AST/ALT in 28 to 67 % cases.

Malaria positive cases and enteric cases were found 13.6 % and 15.2 % respectively in present study which was comparable to a study done by Balvinder et al in which was reported 4.7%. & 10.9% respectively.<sup>[12]</sup> Comorbidities like HTN, DM2, CAD, COPD combined together were present in 9.1% of patients in these studies. Low incidence of comorbidity may be explained by relatively younger people getting affected from the illness, as these comorbidities are usually associated with the advanced age. Most common

age group affected in this study was between 15-45 years of age, reflecting young economically active people are affected more with these illnesses which were also supported by literature.<sup>[13,14]</sup> Increased outdoor exposure to mosquitoes and ticks, use of contaminated water may be factor responsible for this. Males predominate over female in this study. Similar observation was made by other study conducted in northern India.<sup>[15]</sup> It can also be explained with more outdoor activities and more risk of exposure for mosquito bite among males.

Present epidemiological evidence of association of comorbidities with severity of illness is limited and only suggestive. However it is important to identify these comorbidities early and patient should be kept in close observation if needed to improve outcome and avoid complications.<sup>[16]</sup>

#### Limitation of Study

- The limitation of our study is that it excluded the paediatrics age group or else we could have expected to have different fever etiology due to exposure and immunity. Hence, it is prudent that the pattern of fever etiology found in our study is not representative of general population.
- Data to differentiate vivax from falciparum and serial platelet counts could not be collected in this study. Enteric fever cases would have been diagnosed better if tests for blood culture and Widal test data could have been correlated. Though sensitivity and specificity of the serological testing for various aetiologies are not 100% and chances of over or under diagnosis is there. However, this number should not be large enough to interfere with the purpose of the study

#### Conclusion

In our study, predominantly young population (<55 years) was most commonly affected. The most common etiology was found to be Rickettsial Fever followed by Enteric Fever and Malaria. This upholds the necessity for better

health awareness, early transport and presentation along with better diagnostic tools at primary health care facilities. A protocol based approach for diagnosing and managing Acute Febrile illness should be done to reduce the cost, burden on healthcare system.

**Declaration of Conflicts of Interest-** All authors have none to declare.

#### References

1. Blumberg L, Ogunbanjo GA, Durrheim DN. Fever in adults-approach to diagnosis and management, SA Fam Pract .2000; 22:23-26.
2. Joshi R, Colford JM, Reingold A. Nonmalarial acute undifferentiated fever in a rural hospital in central India – Diagnostic uncertainty and overtreatment with antimalarial agents. Am J Trop Med Hyg. 2008;78(3):393-99.
3. Chrispal A, Boorugu H, Gopinath KG, Chandy S, PrakashJA, et al. Acute undifferentiated febrile illness in India. Int Journal of Emer Med. 2011;4:57.
4. Chrispal A, Boorugu H, Gopinath KG, Chandy S, Prakash JA, et al. Acute Undifferentiated febrile illness in adult hospitalized patients: the disease spectrum and diagnostic predictors – an experience from a tertiary care hospital in South India. Trop Doc. 2010;40(4):230-34.
5. Garima Mittal, Sohaib Ahmad, R K Agarwal, Minakshi Dhar, Manish Mittal, et al . Aetiologies of Acute Undifferentiated Febrile illness in Adult Patients – an Experience from a Tertiary Care Hospital in Northern India Journal of Clinical and Diagnostic Research. 2015 Dec, Vol-9(12): DC22-4.
6. Singh R, Singh SP, Ahmad N. A Study of Etiological Pattern in an Epidemic of Acute Febrile Illness during Monsoon in a Tertiary Health Care Institute of

- Uttarakhand, India. *J ClinDiagn Res.* 2014;8(6):MC01-03.
7. Susilawati, T.N., et al. Acute undifferentiated fever in Asia: a review of the literature. *The Southeast Asian J. Trop. Med. Public Health.* 2014, 45(3): 71926.
  8. Manocha, et al. Frequency of leptospirosis in patients of acute febrile illness in Uttar Pradesh. 2004J. *Assoc. Physicians of India*, 52: 623-5.
  9. Phuong, H.L., et al. Acute undifferentiated fever in Binh Thuan Province, Vietnam: Imprecise clinical diagnosis and irrational pharmacotherapy 2006. *Trop. Med. Int. Health*, 11: 869-79
  10. Sharma A, Raina R, Dhiman P, Adarsh, Madhabhavi I, Panda P. Rare Coinfection of Scrub Typhus and Malaria in Immunocompetent Person. *Online J Health Allied Scs.* 2012;11(2):12.
  11. Shashidharan VK, Nagpal AK. Tropical infections in ICU. In: *Handbook of Medical Emergencies*, 2nd Edition, India: Department of Internal Medicine, AFMC Pune; 2015. 306-19.
  12. Balvinder Singh Arora, Monika Matlani, Karnika Saigal. Major aetiologies of acute undifferentiated fever in 2013 and 2014: an experience in retrospect *Int J Adv Med.* 2017 Apr; 4(2):568-572
  13. Sharma A, Raina R, Dhiman P, Adarsh, Madhabhavi I, Panda P. Rare Coinfection of Scrub Typhus and Malaria in Immunocompetent Person. *Online J Health Allied Scs.* 2012; 11(2):12
  14. MA Andrews, Aleena Elizabeth, Praveenlal Kuttichira. Clinical profile of acute undifferentiated febrile illness in patients admitted to a teaching hospital in Kerala. *Health Sciences* 2014;1(3): JS001D.
  15. Garima Mittal, Sohaib Ahmad, R K Agarwal, Minakshi Dhar, Manish Mittal, and Shiwani Sharma. Aetiologies of Acute Undifferentiated Febrile illness in Adult Patients – an Experience from a Tertiary Care Hospital in Northern India. *J ClinDiagn Res.* 2015 Dec; 9(12): 22–24
  16. Pang J, Salim A, Lee VJ, Hibberd ML, Chia KS, Leo YS, et al. Diabetes with hypertension as risk factors for adult dengue haemorrhagic fever in a predominantly dengue serotype 2 epidemic: a case control study. *PLoS Negl Trop Dis.* 2012; 6(5):e1641.