



## Study of Mean Platelet Volume in Acute Coronary Syndrome

Authors

**Prasanth Kumar P<sup>1</sup>, Ramakrishna Rao<sup>2</sup>, Periasamy S<sup>3</sup>**

<sup>1</sup>Post Graduate, Department of General Medicine, Rajah Muthiah Medical College and Hospital, Chidambaram, India, 608002

<sup>2</sup>Professor, Department of General Medicine, Rajah Muthiah Medical College and Hospital, Chidambaram, India, 608002

<sup>3</sup>Professor, Department of General Medicine, Rajah Muthiah Medical College and Hospital, Chidambaram, India, 608002

\*Corresponding Author

**Dr Ramakrishna Rao**

Professor, Department of General Medicine, Rajah Muthiah Medical College and Hospital, Chidambaram, India, 608002

### Abstract

**Background:** As the initial step in pathogenesis of Acute coronary syndrome is plaque erosion, rupture and thrombus formation. Mean platelet volume (MPV) correlates with platelet function and activation. Mean platelet volume can be used as a diagnostic parameter in patients with Acute Coronary syndrome. The primary objective of this study is to find out the association between mean platelet volume and Acute Coronary syndrome.

**Methods:** Our study included 50 cases with acute coronary syndrome and 50 controls who were selected based on inclusion and exclusion criteria. Mean platelet volume estimated by auto-analyser is compared between ST-elevation Myocardial infarction, Non- ST-elevation Myocardial infarction and control group.

**Results:** The mean value of mean platelet volume among those with diagnosis STEMI, NSTEMI and CONTROL was  $11.09 \pm 1.046$  fl,  $10.5 \pm 0.54$ fl, and  $7.84 \pm 0.375$  fl respectively. The mean was found to be statistically significant for each diagnosis.

**Conclusion:** Our study concluded that mean platelet volume is higher in patients with STEMI than NSTEMI but in control group mean platelet volume found to be normal. Hence mean platelet volume can be used as a diagnostic marker in Acute coronary syndrome.

**Keywords:** Acute coronary syndrome, STEMI, NSTEMI, Mean platelet volume.

### Introduction

Acute coronary syndrome consists of unstable angina, STEMI and NSTEMI. Almost all cases of coronary artery disease is caused by atherosclerosis. After rupture of atherosclerotic plaque in the coronary artery, there is platelet adhesion, activation and aggregation of platelets

which leads to formation of thrombus and ultimately resulting in acute myocardial infarction<sup>[1]</sup>. Large platelets are more active and more adhesive and tend to aggregate more than smaller ones<sup>[2]</sup>. Procoagulant properties of large platelets are due to abundance dense granules which contains ADP, P-selectin, Gp IIIa and

serotonin<sup>[3]</sup>. Coarsh et al<sup>[4]</sup> suggested that large platelets are metabolically and enzymatically more active than small platelets as assessed by invitro aggregometry. Mean platelet volume measures average size of platelets in circulating blood. Normal range of mean platelet volume is 7-9 fL<sup>[5]</sup>. The aim of our study is to determine the association between mean platelet volume and acute coronary syndrome

**Methods**

The first 50 patients who were admitted with Acute coronary syndrome based on selection criteria in the department of general medicine in

Rajah Muthiah medical college and hospital were taken for this study. We also took 50 healthy attenders of inpatients with corresponding age and sex who met the exclusion was taken into the study as a control group.

**Inclusion Criteria:** Patients who presented with chest pain lasting 20 minutes or more with CK-MB  $\geq 40$  IU/L or Trop-I positive and ECG showing ST-elevation or ST-depression.

**Exclusion Criteria:** Our study excludes patients with history of Cerebrovascular accidents, Coronary artery disease, Type 2 diabetes mellitus, Systemic hypertension, chronic kidney disease and history of platelet disorders.


**Results**

Total number of study participants enrolled into the study was 100.

**Table 1:** Distribution of study participants according to diagnosis (n=100)

Diagnosis	Frequency	Percentage
STEMI	25	25.0
NSTEMI	25	25.0
CONTROL	50	50.0
Total	100	100

**Fig 1:** Pie chart showing distribution according to diagnosis (n=100)




Out of 100 study participants 50% belonged to control group, 25% with STEMI, and 25% with NSTEMI.

**Table 2:** Distribution of study participants according to age group (n=100)

Age (years)	Frequency	Percentage (%)
<40	8	8.0
41-50	66	66.0
>51	26	26.0
Total	100	100

**Fig 2:** Pie chart showing distribution according to age group (n=100)




Among the 100 study participants 8% were less than 40 years old group, 66% were 41 to 50 years old group, 26% were more than 51 years group.

**Table 3:** Distribution of study participants according to gender (n=100)

Diagnosis	Frequency
Male	61
Female	39

**Fig 3:** Pie chart showing the according to gender (n=100)




Among 100 study participants 61 were males and 39 were females

**Table 4:** Comparison of mean age between STEMI, NSTEMI and CONTROL (n=100)

Diagnosis	Age in years		F value	P value
	Mean	SD		
STEMI	47.84	4.048	.026	.974
NSTEMI	47.96	5.358		
CONTROL	47.70	4.709		

**Fig 4:** Bar chart showing mean age between the groups.




The mean age among those with diagnosis STEMI, NSTEMI and control was  $47.84 \pm 4.048$  years,  $47.96 \pm 5.358$  years and  $47.70 \pm 4.709$  years, respectively. All the three were similar with respect to age.

**Table 5:** Distribution of sex and diagnosis (n=100)

Diagnosis	Male		Female	
	N	%	N	%
STEMI	17	27.9	8	20.5
NSTEMI	15	24.6	10	25.6
CONTROL	29	47.5	21	53.8
Total	61	100	39	100

$\chi^2 - 0.715$   $df - 2$   $P$  value - 0.700

**Fig 5:** Compound bar chart showing distribution of sex and diagnosis




61% of the study participants were males and 39% were females. Among the males, 27.9% had STEMI, 24.6% had NSTEMI. Among the

females, 20.5% had NSTEMI, 25.6% HAD NSTEMI. This association was not statistically significant.

**Table 6:** Comparison of mean platelet count volume between STEMI, NSTEMI, CONTROL (n=100)

Diagnosis	Mean Platelet Volume		F value	P value
	Mean	SD		
STEMI	11.09	1.046	115.539	< 0.001
NSTEMI	10.5	0.54		
CONTROL	7.84	0.375		

**Fig 6:** Bar chart showing mean platelet count volume between the groups




The mean value of mean platelet volume among those with diagnosis STEMI, NSTEMI and CONTROL was  $11.09 \pm 1.046$  fl,  $10.5 \pm 0.54$  fl,

and  $7.84 \pm 0.375$  fl respectively. The mean was found to be statistically significant for each diagnosis.

**Table 7:** Comparison of mean values CK-MB between STEMI, NSTEMI, CONTROL (n=100)

Diagnosis	Mean platelet volume		F value	P value
	Mean	SD		
STEMI	105.60	10.45	224.706	<.001
NSTEMI	83.84	10.119		
CONTROL	22.10	6.482		

**Fig 7:** Bar chart showing mean values CK-MB between the groups



The mean value of mean CK-MB among those with diagnosis STEMI, NSTEMI and CONTROL was  $105.60 \pm 10.45$  IU/L,  $83.84 \pm 10.119$  IU/L and  $22.10 \pm 6.42$  IU/L, respectively. The mean was found to be statistically significant for each diagnosis

**Discussion**

The study done by Randheer et al evaluated 215 patients and results showed that mean platelet volume (MPV) was found to be higher among ACS patients as compared to non-ACS,  $11.44 \pm 1.23$  vs  $9.91 \pm 1.27$  fl ( $p$ -value  $< 0.001$ )<sup>[6]</sup>. In another study done by pervin et al, showed results similar to our study (MPV higher in ACS than Non-ACS)<sup>[7]</sup>. The study done by A Mathur et al and AS Assiri et al are also in concordance with this study<sup>[8,9]</sup>. This study also compares mean platelet volume among STEMI and NSTEMI case groups. The result showed that mean value of MPV in STEMI is  $10.32 \pm 0.77932$  which was greater than the mean value of MPV in NSTEMI which was  $9.22 \pm 0.52743$  and It was statistically significant ( $P < 0.05$ ). Other studies comparing the MPV in STEMI and NSTEMI was conducted by Rifat et al and the study showed that the difference in mean platelet volume in STEMI and

NSTEMI was  $8.7 \pm 1$  and  $7.9 \pm 0.7$  fl ( $p < 0.01$ ) respectively and it is statistically significant and concordance with this study<sup>[10]</sup>.

**Conclusion**

In this study mean platelet volume was higher in ACS than in control group and in particular mean platelet volume was higher among patients with STEMI than NSTEMI (case groups). Large platelets are more active haemostatically and leads to development of coronary thrombosis. Hence this study showed that mean platelet volume might be useful as an additional diagnostic tool in addition to other diagnostic marker on early detection of myocardial infarction.

**Conflict of interest:** None to declare

**References**

1. Endler G, Klimesch A, Sunder-Plassmann H, Schillinger M, Exner M, Mannhalter C, et al. Mean platelet volume is an independent risk factor for myocardial infarction but not for coronary artery disease. Br. J. Haematol., 2002

2. Furie B, Furie BC. Mechanism of thrombus formation. *N Engl J med* 2008;359(9):938-49.
3. Thompson CB, Jakubowski JA, Quinn PG, et al. Platelet size as a determinant of platelet function. *J Lab Clin Med* 1981;101(2):205-213.
4. Corash L. The relationship between megakaryocyte ploidy and platelet volume. *Blood cells* 1989 ;15:81.
5. Paulus JM. platelet size in man. *Blood* 1975;46:321
6. Pal R, Bagarhatta R, Gulati S, Rathore M, Sharma N. Mean platelet volume in acute coronary syndrome. *JCDR*. 2014;8(8):1-4.
7. Pervin S, Ferdousy S, Hossain M, Joarder AI, Sultana T. Elevated mean platelet volume is a marker of acute coronary syndrome. *Bangladesh Med J*. 2013;42 (2);45-50.
8. Mathur A, Robinson MS, Cotton J, Martin JF, Erusalimsky JD. Platelet reactivity in acute coronary syndromes: evidence for differences in platelet behaviour between unstable angina and myocardial infarction. *Thrombosis and Haemostasis*. 2001;85: 989-94.
9. Assiri AS, Jamil AM, Mahfouz AA, Mahmoud AS, Ghallab S. Diagnostic importance of platelet parameters in patients with acute coronary syndrome admitted to a tertiary care hospital in southwest region, Saudi Arabia, *J Saudi Heart Assoc*. 2012;24(1):17-21.
10. Ulusoy RE, Yokuşoğlu M, Kırılmaz A, Nevruz O, Baysan O, Kılıçaslan F, et al. Mean platelet volume in ST elevation and non-ST elevation myocardial infarction. *Gulhane Med J*. 2011; 53(2):114-8.